

Technical Field

10

15

20

Furthermore, the present inventors has revealed that the *rhtA* gene exists at 18 min on *E. coli*

chromosome and that the *rhtA* gene is identical to
ORF1 between *pexB* and *ompX* genes. The unit
expressing a protein encoded by the ORF1 has been
designated as *rhtA* (*rht*: resistance to homoserine and
5 threonine) gene. The *rhtA* gene includes a 5'-
noncoding region including SD sequence, ORF1 and a
terminator. Also, the present inventors have found
that a wild type *rhtA* gene participates in resistance
to threonine and homoserine if cloned in a multicopy
10 state and that enhancement of expression of the *rhtA*
gene improves amino acid productivity of a bacterium
belonging to the genus *Escherichia* having an ability
to produce L-lysine, L-valine or L-threonine
(ABSTRACTS of 17th International Congress of
15 Biochemistry and Molecular Biology in conjugation
with 1997 Annual Meeting of the American Society for
Biochemistry and Molecular Biology, San Francisco,
California August 24-29, 1997, abstract No. 457).

It is found that at least two different genes
20 which impart homoserine resistance in a multicopy
state exist in *E. coli* during cloning of the *rhtA*
gene. One of the genes is the *rhtA* gene, however the
other gene has not been elucidated.

Disclosure of the Invention

25 An object of the present invention is to provide

09847392-050301
"2624860"

a novel gene participating in resistance to homoserine, and a method for producing an amino acid, especially, L-homoserine, L-alanine, L-isoleucine, L-valine and L-threonine with a high yield.

5 The inventors have found that a region at 86 min on *E. coli* chromosome, when cloned by a multicopy vector, impart resistance to L-homoserine to cells of *E. coli*, and that when the region is amplified, the amino acid productivity of *E. coli* can be improved
10 like the *rhtA* gene. On the basis of these findings, the present invention have completed.

 Thus, the present invention provides:

(1) a DNA coding for a protein as defined in the following (A) or (B):

15 (A) a protein which comprises an amino acid sequence shown in SEQ ID NO: 2 in Sequence Listing;
 or

 (B) a protein which comprises an amino acid sequence including deletion, substitution, insertion
20 or addition of one or several amino acids in the amino acid sequence shown in SEQ ID NO: 2 in Sequence Listing, and which has an activity of making a bacterium having the protein L-homoserine-resistant,

(2) the DNA according to (1), which is a DNA as
25 defined in the following (a) or (b):

(a) a DNA which comprises a nucleotide sequence

05047392-050301
FOE060" 26274860

corresponding to the nucleotide numbers of 557 to 1171 of a nucleotide sequence shown in SEQ ID NO: 1 in Sequence Listing; or

(b) a DNA which hybridizes with the nucleotide sequence corresponding to the nucleotide numbers of 557 to 1171 of the nucleotide sequence shown in SEQ ID NO: 1 in Sequence Listing under stringent conditions, and which codes for the protein having the activity of making the bacterium having the protein L-homoserine-resistant,

(3) a bacterium belonging to the genus *Escherichia*, wherein L-homoserine resistance of the bacterium is enhanced by amplifying a copy number of the DNA of (1) in a cell of the bacterium,

(4) the bacterium of (3), wherein the DNA of (1) is carried on a multicopy vector in the cell of the bacterium,

(5) the bacterium of (3), wherein the DNA of (1) is carried on a transposon in the cell of the bacterium,

(6) a method for producing an amino acid, comprising the steps of cultivating the bacterium of any of (3) to (5), which has an ability to produce the amino acid, in a culture medium to produce and accumulate the amino acid in the medium, and recovering the amino acid from the medium, and

(7) the method of (6), wherein the amino acid is at least one selected from the group consisting of L-

05001-2624860

homoserine, L-alanine, L-isoleucine, L-valine and L-threonine.

5 The DNA of the present invention may be referred to as "*rhtB* gene", a protein coded by the *rhtB* gene may be referred to as "RhtB protein", an activity of the RhtB protein which participates in resistance to L-homoserine of a bacterium (i.e. an activity of making a bacterium having the RhtB protein L-homoserine-resistant) may be referred to as "Rh
10 activity", and a structural gene encoding the RhtB protein in the *rhtB* gene may be referred to as "*rhtB* structural gene". The term "enhancing the Rh activity" means imparting resistance to homoserine to a bacterium or enhance the resistance by means of
15 increasing the number of molecules of the RhtB protein, increasing a specific activity of the RhtB protein, or desensitizing negative regulation against the expression or the activity of the RhtB protein or the like. The terms "DNA coding for a protein" mean
20 a DNA of which one of strands codes for the protein when the DNA is double-stranded. The L-homoserine resistance means a property that a bacterium grows on a minimal medium containing L-homoserine at a concentration at which a wild type strain thereof can
25 not grow, usually at 10 mg/ml. The ability to produce an amino acid means a property that a

09847397.050301
"2624860"

bacterium produces and accumulates the amino acid in a medium in a larger amount than a wild type strain thereof.

According to the present invention, resistance to homoserine of a high concentration can be imparted to a bacterium belonging to the genus *Escherichia*. A bacterium belonging to the genus *Escherichia*, which has increased resistance homoserine and an ability to accumulate an amino acid, especially, L-homoserine, L-alanine, L-isoleucine, L-valine or L-threonine in a medium with a high yield.

The present invention will be explained in detail below.

<1> DNA of the present invention

The DNA of the present invention coding for a protein having the Rh activity and having an amino acid sequence shown in SEQ ID NO: 2 in Sequence Listing. Specifically, the DNA of the present invention may be exemplified by a DNA comprising a nucleotide sequence of the nucleotide numbers 557 to 1171 of a nucleotide sequence shown in SEQ ID NO: 1 in Sequence Listing.

The DNA of the present invention includes a DNA fragment encoding the RhtB protein conferring bacterium *Escherichia coli* resistance to homoserine, which includes the regulatory elements of the *rhtB*

09847299-050301
1000050-25004860

gene and the structural part of *rhtB* gene, having the nucleotide sequence shown in SEQ ID NO: 1.

5 The nucleotide sequence shown in SEQ ID NO: 1 corresponds to a part of sequence complement to the sequence of GenBank accession number M87049. SEQ ID NO: 1 includes f138 (nucleotide numbers 61959-61543 of GenBank accession number M87049) which is a known but function-unknown ORF (open reading frame) present at 86 min on *E. coli* chromosome, and 5'-flanking and 10 3'-flanking regions thereof. The f138, which had only 160 nucleotides in the 5'-flanking region, could not impart the resistance to homoserine. No termination codon is present between the 62160 and 61959 of M87049 (upstream the ORF f138). Hence, the 15 coding region is 201 bp longer. Thus the RhtB protein and the *rhtB* gene coding for the protein are novel.

20 The *rhtB* gene may be obtained, for example, by infecting Mucts lysogenic strain of *E. coli* using a lysate of a lysogenic strain of *E. coli* such as K12 or W3110 according to the method in which mini-Mu d5005 phagemid is used (Groisman, E. A., et al., J. Bacteriol., 168, 357-364 (1986)), and isolating 25 plasmid DNAs from colonies growing on a minimal medium containing kanamycin (40 μ g/ml) and L-homoserine (10 mg/ml). As illustrated in the Example

described below, the *rhtB* gene was mapped at 86 min on the chromosome of *E. coli*. Therefore, the DNA fragment including the *rhtB* gene may be obtained from the chromosome of *E. coli* by colony hybridization or PCR (polymerase chain reaction, refer to White, T.J. et al, Trends Genet. 5, 185(1989)) using oligonucleotide(s) which has a sequence corresponding to the region near the portion of 86 min on the chromosome of *E. coli*. Alternatively, the oligonucleotide may be designed according to the nucleotide sequence shown in SEQ ID NO: 1. By using oligonucleotides having nucleotide sequences corresponding to a upstream region from the nucleotide number 557 and a downstream region from the nucleotide number 1171 in SEQ ID NO: 1 as the primers for PCR, the entire coding region can be amplified.

Synthesis of the oligonucleotides can be performed by an ordinary method such as a phosphoamidite method (see Tetrahedron Letters, 22, 1859 (1981)) by using a commercially available DNA synthesizer (for example, DNA Synthesizer Model 380B produced by Applied Biosystems). Further, the PCR can be performed by using a commercially available PCR apparatus (for example, DNA Thermal Cycler Model PJ2000 produced by Takara Shuzo Co., Ltd.), using *Taq*

DNA polymerase (supplied by Takara Shuzo Co., Ltd.)
in accordance with a method designated by the
supplier.

5 The DNA coding for the RhtB protein of the
present invention may code for RhtB protein including
deletion, substitution, insertion, or addition of one
or several amino acids at one or a plurality of
positions, provided that the Rh activity of RhtB
protein encoded thereby is not deteriorated. The
10 DNA, which codes for the substantially same protein
as the RhtB protein as described above, may be
obtained, for example, by modifying the nucleotide
sequence, for example, by means of the site-directed
mutagenesis method so that one or more amino acid
15 residues at a specified site involve deletion,
substitution, insertion or addition. DNA modified as
described above may be obtained by the conventionally
known mutation treatment. The mutation treatment
includes a method for treating a DNA coding for the
20 RhtB protein *in vitro*, for example, with
hydroxylamine, and a method for treating a
microorganism, for example, a bacterium belonging to
the genus *Escherichia* harboring a DNA coding for the
RhtB protein with ultraviolet irradiation or a
25 mutating agent such as N-methyl-N'-nitro-N-
nitrosoguanidine (NTG) and nitrous acid usually used

for the mutation treatment.

5 The DNA, which codes for substantially the same
protein as the RhtB protein, can be obtained by
expressing a DNA subjected to in vitro mutation
treatment as described above in multicopy in an
appropriate cell, investigating the resistance to
homoserine, and selecting the DNA which increase the
resistance. Also, it is generally known that an
amino acid sequence of a protein and a nucleotide
10 sequence coding for it may be slightly different
between species, strains, mutants or variants, and
therefore the DNA, which codes for substantially the
same protein, can be obtained from L-homoserine-
resistant species, strains, mutants and variants
15 belonging to the genus *Escherichia*. Specifically,
the DNA, which codes for substantially the same
protein as the RhtB protein, can be obtained by
isolating a DNA which hybridizes with DNA having, for
example, a nucleotide sequence of the nucleotide
20 numbers 557 to 1171 of the nucleotide sequence shown
in SEQ ID NO: 1 in Sequence Listing under stringent
conditions, and which codes for a protein having the
Rh activity, from a bacterium belonging to the genus
Escherichia which is subjected to mutation treatment,
25 or a spontaneous mutant or a variant of a bacterium
belonging to the genus *Escherichia*. The term

09847392.050301

"stringent conditions" referred to herein is a condition under which so-called specific hybrid is formed, and non-specific hybrid is not formed. It is difficult to clearly express this condition by using any numerical value. However, for example, the stringent conditions include a condition under which DNAs having high homology, for example, DNAs having homology of not less than 70% with each other are hybridized, and DNAs having homology lower than the above with each other are not hybridized.

<2> Bacterium belonging to the genus *Escherichia* of the present invention

The bacterium belonging the genus *Escherichia* of the present invention is a bacterium belonging to the genus *Escherichia* of which the Rh activity is enhanced. A bacterium belonging to the genus *Escherichia* is exemplified by *Escherichia coli*. The Rh activity can be enhanced by, for example, amplification of the copy number of the *rhtB* structural gene in a cell, or transformation of a bacterium belonging to the genus *Escherichia* with a recombinant DNA in which a DNA fragment including the *rhtB* structural gene encoding the RhtB protein is ligated with a promoter sequence which functions efficiently in a bacterium belonging to the genus *Escherichia*. The Rh activity can be also enhanced by

substitution of the promoter sequence of the *rhtB* gene on a chromosome with a promoter sequence which functions efficiently in a bacterium belonging to the genus *Escherichia*.

5 The amplification of the copy number of the *rhtB* structural gene in a cell can be performed by introduction of a multicopy vector which carries the *rhtB* structural gene into a cell of a bacterium belonging to the genus *Escherichia*. Specifically,
10 the copy number can be increased by introduction of a plasmid, a phage or a transposon (Berg, D.E. and Berg, C. M., Bio/Technol., 1, 417 (1983)) which carries the *rhtB* structural gene into a cell of a bacterium belonging to the genus *Escherichia*.

15 The multicopy vector is exemplified by plasmid vectors such as pBR322, pMW118, pUC19 or the like, and phage vectors such as λ 1059, λ BF101, M13mp9 or the like. The transposon is exemplified by Mu, Tn10, Tn5 or the like.

20 The introduction of a DNA into a bacterium belonging to the genus *Escherichia* can be performed, for example, by a method of D. M. Morrison (Methods in Enzymology 68, 326 (1979)) or a method in which recipient bacterial cells are treated with calcium
25 chloride to increase permeability of DNA (Mandel, M. and Higa, A., J. Mol. Biol., 53, 159 (1970)) and the

like.

If the Rh activity is enhanced in an amino acid-producing bacterium belonging to the genus *Escherichia* as described above, a produced amount of the amino acid can be increased. As the bacterium belonging to the genus *Escherichia* which is to be the Rh activity is enhanced, strains which have abilities to produce desired amino acids are used. Besides, an ability to produce an amino acid may be imparted to a bacterium in which the Rh activity is enhanced. Examples of amino acid-producing bacteria belonging to the genus *Escherichia* are described below.

(1) L-threonine-producing bacteria

The L-threonine-producing bacteria belonging to the genus *Escherichia* may be exemplified by strain MG442 (Guayatiner et al., Genetika (in Russian), 14, 947-956 (1978)).

(2) L-homoserine-producing bacteria

The L-homoserine-producing bacteria belonging to the genus *Escherichia* may be exemplified by strain NZ10 (*thrB*). This strain was derived from the known strain C600 (*thrB*, *leuB*) (Appleyard R.K., Genetics, 39, 440-452 (1954)) as Leu⁺ revertant.

On the basis of the *rhtB* DNA fragment, new amino acid-producing strains *E. coli* NZ10/pAL4,pRhtB; *E. coli* MG422/pVIC40,pRhtB; and *E. coli* MG442/pRhtB were

obtained which are used for the production of amino acids by fermentation.

5 The new strains have been deposited (according to international deposition based on Budapest Treaty) in the Russian National Collection of Industrial Microorganisms (VKPM) on October 6, 1998. The strain *E. coli* NZ10/pAL4,pRhtB has been deposited as an accession number of VKPM B-7658; the strain *E. coli* MG442/pRhtB has been deposited as an accession number of VKPM B-7659; and the strain *E. coli* MG442/pVIC40,pRhtB has been deposited as an accession number of VKPM B-7660.

15 The strain *E. coli* NZ10/pAL4,pRhtB (VKPM B-7658) exhibits the following cultural-morphological and biochemical features.

Cytomorphology. Gram-negative weakly-motile rods having rounded ends. Longitudinal size, 1.5 to 2 μ m.

Cultural features:

20 Beef-extract agar. After 24-hour growth at 37°C, produces round whitish semitransparent colonies 1.5 to 3 mm in diameter, featuring a smooth surface, regular or slightly wavy edges, the center is slightly raised, homogeneous structure, pastelike consistency, readily emulsifiable.

25 Luria's agar. After a 24-hour growth at 37°C, develops whitish semitranslucent colonies 1.5 to 2.5

09847391-050301

mm in diameter having a smooth surface, homogeneous structure, pastelike consistency, readily emulsifiable.

Minimal agar-doped medium M9. After 40 to 48 hours of growth at 37°C, forms colonies 0.5 to 1.5 mm in diameter, which are colored greyish-white, semitransparent, slightly convex, with a lustrous surface.

Growth in a beef-extract broth. After 24-hour growth at 37°C, exhibits strong uniform cloudiness, has a characteristic odor.

Physiological and biochemical features:

Grows upon thrust inoculation in a beef-extract agar. Exhibits good growth throughout the inoculated area.

The microorganism proves to be a facultative anaerobe.

It does not liquefy gelatin.

Features a good growth on milk, accompanied by milk coagulation.

Does not produce indole.

Temperature conditions. Grows on beef-extract broth at 20-42°C, an optimum temperature lying within 33-37°C.

pH value of culture medium. Grows on liquid media having the pH value from 6 to 8, an optimum value being 7.2.

Carbon sources. Exhibits good growth on glucose,

09047392 050301
T0E050 26E24860

fructose, lactose, mannose, galactose, xylose, glycerol, and mannitol to produce an acid and gas. Nitrogen sources. Assimilates nitrogen in the form of ammonium, nitric acid salts, as well as from some organic compounds.

Resistant to ampicillin, kanamycin and L-homoserine. L-Threonine is used as a growth factor.

Content of plasmids. The cells contain multicopy hybrid plasmid pAL4 ensuring resistance to ampicillin and carrying the gene *thrA* of the threonine operon, which codes for aspartate kinase-homoserine dehydrogenase I responsible for the increased homoserine biosynthesis. Besides, the cells contain a multicopy hybrid plasmid pRhtB ensuring resistance to kanamycin and carrying the *rhtB* gene which confers resistance to homoserine (10 mg/l).

The strain *E. coli* MG442/pRhtB (VKPM B-7659) has the same cultural-morphological and biochemical features as the strain NZ10/pAL4,pRhtB except for L-isoleucine is used as a growth factor instead of L-threonine. However, the strain can grow slowly without isoleucine. Besides, the cells of the strain contain only one multicopy hybrid plasmid pRhtB ensuring resistance to kanamycin and carrying the *rhtB* gene which confers resistance to homoserine (10 mg/l).

The strain *E. coli* MG442/pVIC40,pRhtB (VKPM B-7660) has the same cultural-morphological and biochemical features as the strain NZ10/pAL4,pRhtB except for L-isoleucine is used as a growth factor instead of L-threonine. However, the strain can grow slowly without isoleucine. The cells of the strain contain multicopy hybrid plasmid pVIC40 ensuring resistance to streptomycin and carrying the genes of the threonine operon. Besides, they contain multicopy hybrid plasmid pRhtB ensuring resistance to kanamycin and carrying the *rhtB* gene which confers resistance to homoserine (10 mg/l).

<3> Method for producing an amino acid

An amino acid can be efficiently produced by cultivating the bacterium in which the Rh activity is enhanced by amplifying a copy number of the *rhtB* gene as described above, and which has an ability to produce the amino acid, in a culture medium, producing and accumulating the amino acid in the medium, and recovering the amino acid from the medium. The amino acid is exemplified preferably by L-homoserine, L-alanine, L-isoleucine, L-valine and L-threonine.

In the method of present invention, the cultivation of the bacterium belonging to the genus *Escherichia*, the collection and purification of amino

acid from the liquid medium may be performed in a manner similar to those of the conventional method for producing an amino acid by fermentation using a bacterium. A medium used in cultivation may be
5 either a synthetic medium or a natural medium, so long as the medium includes a carbon and a nitrogen source and minerals and, if necessary, nutrients which the bacterium used requires for growth in appropriate amounts. The carbon source may include
10 various carbohydrates such as glucose and sucrose, and various organic acids. Depending on assimilatory ability of the used bacterium, alcohol including ethanol and glycerol may be used. As the nitrogen source, ammonia, various ammonium salts such as
15 ammonium sulfate, other nitrogen compounds such as amines, a natural nitrogen source such as peptone, soybean hydrolyte and digested fermentative microbe are used. As minerals, monopotassium phosphate, magnesium sulfate, sodium chloride, ferrous sulfate,
20 manganese sulfate, calcium carbonate are used.

The cultivation is preferably culture under an aerobic condition such as a shaking culture, and an aeration and stirring culture. The temperature of culture is usually 20 to 40°C, preferably 30 to 38°C.
25 The pH of the culture is usually between 5 and 9, preferably between 6.5 and 7.2. The pH of the culture can be adjusted with ammonia, calcium

09047392 050304

carbonate, various acids, various bases, and buffers. Usually, a 1 to 3-day cultivation leads to the accumulation of the target amino acid in the medium.

Recovering the amino acid can be performed by removing solids such as cells from the medium by centrifugation or membrane filtration after cultivation, and then collecting and purifying the target amino acid by ion exchange, concentration and crystalline fraction methods and the like.

Brief Explanation of Drawing

Fig. 1 shows cloning, identification and inactivation of the *rhtB* gene.

Fig. 2 shows the amino acid sequence of the RhtB protein.

Examples

The present invention will be more concretely explained below with reference to Examples. In the Examples, an amino acid is of L-configuration unless otherwise noted.

Example 1: Obtaining of *rhtB* DNA fragment

(1) Cloning of *rhtB* gene into mini-Mu phagemid

The wild-type *rhtB* gene was cloned *in vivo* using mini-Mu d5005 phagemid (Groisman, E. A., et al., J.

Bacteriol., 168, 357-364 (1986)). MuCts62 lysogen of the strain MG442 was used as a donor. Freshly prepared lysates were used to infect a Mucts lysogenic derivative of a strain VKPM B-513 (Hfr K10 metB). The cells were plated on M9 glucose minimal medium with methionine (50 µg/ml), kanamycin (40 µg/ml) and homoserine (10 mg/ml). Colonies which appeared after 48 hr were picked and isolated. Plasmid DNA was isolated and used to transform the strain VKPM B-513 by standard techniques. Transformants were selected on L-broth agar plates with kanamycin as above. Plasmid DNA was isolated from those which were resistant to homoserine, and analyzed by restriction mapping of the structure of the inserted fragments. It appeared that two types of inserts belonging to different chromosome regions had been cloned from the donor. Thus, at least two different genes that is in multicopy and imparts resistance to homoserine exist in *E. coli*. One of the two type of inserts is the *rhtA* gene which has already reported (ABSTRACTS of 17th International Congress of Biochemistry and Molecular Biology in conjugation with 1997 Annual Meeting of the American Society for Biochemistry and Molecular Biology, San Francisco, California August 24-29, 1997). Among the other of the two types of inserts, a fragment of a

09847399-050304

minimum length which imparts the resistance to
homoserine is of 0.8 kb (Fig. 1).

(2) Identification of *rhtB* gene

5 The insert fragment was sequenced by the dideoxy
chain termination method of Sanger. Both DNA strands
were sequenced in their entirety and all junctions
were overlapped. The sequencing showed that the
insert fragment included f138 (nucleotide numbers
61543 to 61959 of GenBank accession number M87049)
10 which was a known but function-unknown ORF (open
reading frame) present at 86 min of *E. coli*
chromosome and 201 bp of an upstream region thereof
(downstream region in the sequence of M87049). The
f138 which had only 160 nucleotides in the 5'-
15 flanking region could not impart the resistance to
homoserine. No termination codon is present upstream
the ORF f138 between 62160 and 61959 nucleotides of
M87049. Furthermore, one ATG following a sequence
predicted as a ribosome binding site is present in
20 the sequence. The larger ORF (nucleotide numbers
62160 to 61546) is designated as *rhtB* gene. The RhtB
protein deduced from the gene is highly hydrophobic
and contains 5 possible transmembrane segments.

Example 2: Production of homoserine-producing strain

25 Strain NZ10 of *E. coli* was transformed by a

plasmid pAL4 which was a pBR322 vector into which the
thrA gene coding for aspartokinase-homoserine
dehydrogenase I was inserted, to obtain the strains
NZ10/pAL4. The strain NZ10 is a *leuB*⁺-reverted mutant
5 (*thrB*) obtained from the *E. coli* strain C600 (*thrB*,
leuB) (Appleyard, Genetics, 39, 440-452 (1954)).

The *rhtB* gene was inserted to a plasmid pUK21
which is a known plasmid pUC19 in which a kanamycin
resistance gene substituted for an ampicillin
10 resistance gene (Vieira, J. and Messing, J., Gene,
100, 189-194 (1991)), to obtain pRhtB.

The strain NZ10/pAL4 was transformed with pUK21
or pRhtB to obtain strains NZ10/pAL4,pUK21 and
NZ10/pAL4,pRhtB.

15 The thus obtained transformants were each
cultivated at 37°C for 18 hours in a nutrient broth
with 50 mg/l kanamycin and 100 mg/l ampicillin, and
0.3 ml of the obtained culture was inoculated into 3
ml of a fermentation medium having the following
20 composition and containing 50 mg/l kanamycin and 100
mg/l ampicillin, in a 20 x 200 mm test tube, and
cultivated at 37°C for 46 hours with a rotary shaker.
After the cultivation, an accumulated amount of
homoserine in the medium and an absorbance at 560 nm
25 of the medium were determined by known methods.

09047399-050301
"26E24860"

Fermentation medium composition (g/L)

	Glucose	80
	(NH ₄) ₂ SO ₄	22
	K ₂ HPO ₄	2
5	NaCl	0.8
	MgSO ₄ ·7H ₂ O	0.8
	FeSO ₄ ·7H ₂ O	0.02
	MnSO ₄ ·5H ₂ O	0.02
	Thiamine hydrochloride	0.0002
10	Yeast Extract	1.0
	CaCO ₃	30
	(CaCO ₃ was separately sterilized.)	

The results are shown in Table 1. As shown in Table 1, the strain NZ10/pAL4,pRhtB accumulated homoserine in a larger amount than the strains NZ10/pAL4 and NZ10/pAL4,pUK21 in which the *rhtB* gene was not enhanced.

Table 1

Strain	OD ₅₆₀	Accumulated amount of homoserine (g/L)
NZ10/pAL4	16.4	3.1
NZ10/pAL4,pUK21	14.3	3.3
NZ10/pAL4,pRhtB	15.6	6.4

Example 3: Production of alanine, valine and isoleucine
with pRhtB-introduced strain

E. coli strain MG442 is a known strain (Gusyatiner, et al., 1978, Genetika (in Russian), 14:947-956).

5 The strain MG442 was transformed with the plasmids pUK21 and pRhtB to obtain strains MG442/pUK21 and MG442/pRhtB.

10 The thus obtained transformants were each cultivated at 37°C for 18 hours in a nutrient broth with 50 mg/l kanamycin, and 0.3 ml of the obtained culture was inoculated into 3 ml of the fermentation medium described in Example 3 and containing 50 mg/l kanamycin, in a 20 x 200 mm test tube, and cultivated at 37°C for 40 hours with a rotary shaker. After the
15 cultivation, accumulated amounts of alanine, valine and isoleucine in the medium and an absorbance at 560 nm of the medium were determined by known methods.

20 The results are shown in Table 2. As shown in Table 2, the strain MG442/pRhtB accumulated each of alanine, valine and isoleucine in a larger amount than the strain MG442/pUK21 in which the *rhtB* gene was not enhanced.

Table 2

Strain	OD ₅₆₀	Accumulated amount (g/L)		
		Alanine	Valine	Isoleucine
MG442/pUK21	13.4	0.2	0.2	0.3
MG442/pRhtB	13.7	0.7	0.5	0.5

Example 4: Production of threonine-producing strain

The strain MG442 (Example 3) was transformed by introducing a known plasmid pVIC40 (U.S. Patent No. 5,175,107 (1992)) by an ordinary transformation method. Transformants were selected on LB agar plates containing 0.1 mg/ml streptomycin. Thus a novel strain MG422/pVIC40 was obtained.

The strain MG442/pVIC40 was transformed with pUK21 or pRhtB to obtain strains MG442/pVIC40,pUK21 and MG442/pVIC40,pRhtB.

The thus obtained transformants were each cultivated at 37°C for 18 hours in a nutrient broth with 50 mg/l kanamycin and 100 mg/l streptomycin, and 0.3 ml of the obtained culture was inoculated into 3 ml of the fermentation medium described in Example 3 and containing 50 mg/l kanamycin and 100 mg/l streptomycin, in a 20 x 200 mm test tube, and cultivated at 37°C for 46 hours with a rotary shaker. After the cultivation, an accumulated amount of threonine in the medium and an absorbance at 560 nm of the medium were determined by

known methods.

The results are shown in Table 3. As shown in Table 3, the strain MG442/pVIC40,pRhtB accumulated threonine in a larger amount than the strains
5 MG442/pVIC40 and MG442/pVIC40,pUK21 in which the *rhtB* gene was not enhanced.

Table 3

Strain	OD ₅₆₀	Accumulated amount of threonine (g/L)
MG442/pVIC40	17	13.6
MG442/pVIC40,pUK21	16.3	12.9
MG442/pVIC40,pRhtB	15.2	16.3

Example 5: Effect of *rhtB* gene inactivation and amplification on bacterium *E. coli* resistance to some amino acids and amino acid analogues

To inactivate the chromosomal *rhtB* gene the plasmid
5 pNPZ46 was constructed (Fig. 1) on the basis of pUK21 vector. It harbors a DNA fragment from 86 min of *E. coli* chromosome, with the *rhtB* gene and 5'-flanking and 3'-flanking regions thereof. Then the *Cla*I-*Eco*47III fragment of the pNPZ46 plasmid *rhtB* gene was
10 substituted for *Asu*II-*Bsr*BI fragment containing *cat* (*Cm*^R) gene of pACYC184 plasmid (Chang and Cohen, J. Bacteriol., 134, 1141-1156, 1978) giving the pNPZ47

plasmid (Fig. 1). To introduce the obtained
insertionally inactivated *rhtB* gene into the chromosome
of the *E. coli* strain N99 (the streptomycin-resistant
derivative of the known strain W3350 (Campbell,
5 Virology, 14, 22-33, 1961)), the method of Parker and
Marinus was used (Parker, B. and Marinus, M. G., Gene,
73, 531-535, 1988). The substitution of the wild type
allele for the inactivated one was proved by phage P1
transduction and by Southern hybridization (Southern,
10 E. M., J. Mol. Biol., 98, 503-517, 1975).

Then the susceptibility of the thus obtained *E.*
coli strain N99 *rhtB::cat*, of the initial strain N99
(*rhtB*⁻) and of its derivative transformed with pRhtB
plasmid, N99/pRhtB, to some amino acids and amino acid
15 analogues was tested. Overnight cultures of the
strains grown in M9 minimal medium at 37°C with a
rotary shaker (10⁹ cfu/ml) were diluted 1:100 and grown
for 5 hours under the same conditions. Then the log
phase cultures thus obtained were diluted and about 10⁴
20 of alive cells were applied to well-dried test plates
with M9 agar containing doubling increments of amino
acids or analogues. The minimum inhibitory
concentration (MIC) of these compounds were examined
after 40-46 h cultivation. The results are shown in
25 Table 4.

Table 4

Substrate	MIC ($\mu\text{g/ml}$)		
	N99(<i>rhtB</i> ⁺)	N99/pRhtB	N99 <i>rhtB::cat</i>
1. L-homoserine	250	30000	125
2. L-threonine	30000	50000	30000
3. L-serine	5000	10000	5000
4. L-valine	0.5	1.	0.5
5. AHVA	50	2000	25
6. AEC	10	25	10
7. 4-aza-DL-leucine	40	100	40

It follows from the Table 4 that multiple copies of *rhtB* besides homoserine conferred upon cells increased resistance to threonine, serine, valine, α -amino- β -hydroxyvaleric-acid (AHVA), S-(2-aminoethyl)-L-cysteine (AEC), and 4-aza-DL-leucine. The inactivation of the *rhtB* gene, on the contrary, increased the cell sensitivity to homoserine and AHVA. These results in conjunction with the data on homology of the RhtB protein to LysE lysine efflux transporter of *Corynebacterium glutamicum* (Vrljic et al., Mol. Microbiol., 22, 815-826, 1996) indicate the analogues function for the *rhtB* gene product. The presumed efflux transporters, RhtB, has specificity to several substrates (amino acids), or may show non-specific effects as a result of amplification.

SEQUENCE LISTING

<110> Livshits, Vitaly Arkadievich
 Zakataeva, Natalya Pavlovna
 Alcoshin, Vladimir Venyamiovich
 Belarcova, Alla Valentinovna
 Tokhmakova, Irina Lvovna

<120> DNA coding for protein which confers on bacterium Escherichia coli
 resistance to L-homoserine, and method for producing L-amino acids

<130>

<140> RU 98118425

<141> 1998-10-13

<160> 2

<170> PatentIn Ver. 2.0

<210> 1

<211> 1200

<212> DNA

<213> Escherichia coli

<220>

<221> CDS

<222> (557)..(1171)

<400> 1

agaaataatg tggagatcgc accgcccatac gaatgtgcca gtatatagcg tttacgccac 60
 ggaccgggct gaacctcctg ctgccagaat gccgccagat catcaacata atcattaaag 120
 cgattaacat gcccgagatg cggatcgggt aacaggcgac cggaacgtcc ctgcccgcga 180
 tggctgatga ttaagacatc aaaccccaaa tggaacaggt cataggccag ttccgcatat 240
 tttacgtagc tctcaatacg ccccgggcag atgactacca cccggtcatg gtgctgtgcg 300
 cgaaaacgga caaagcgcac cggaatgtca tccacaccag taaactctgc ttcacacgc 360
 tgacgccaga aatcagtcag cgggtcccatg gtaaaagcag caaacgcggt ttctcttgtt 420
 tcccagtcct tttgctgctg aaacatcggg taatctgcct cttaaaccac gtaaaatcgt 480
 ttttttttagc gtgcctgaca caacgtcgcg acagtagcgt attgtggcac aaaaatagac 540
 acaccgggag ttcatac atg acc tta gaa tgg tgg ttt gcc tac ctg ctg aca 592
 Met Thr Leu Glu Trp Trp Phe Ala Tyr Leu Leu Thr
 1 5 10
 tcg atc att tta acg ctg tcg cca ggc tct ggt gca atc aac act atg 640
 Ser Ile Ile Leu Thr Leu Ser Pro Gly Ser Gly Ala Ile Asn Thr Met
 15 20 25
 acc acc tcg ctc aac cac ggt tat ccg gcc ggt ggc gtc tat tgc tgg 688
 Thr Thr Ser Leu Asn His Gly Tyr Pro Ala Gly Gly Val Tyr Cys Trp
 30 35 40
 gct tca gac cgg act ggc gat tca tat tgt gct ggt tgg cgt ggg gtt 736
 Ala Ser Asp Arg Thr Gly Asp Ser Tyr Cys Ala Gly Trp Arg Gly Val
 45 50 55 60
 ggg acg cta ttt tcc cgc tca gtg att gcg ttt gaa gtg ttg aag tgg 784
 Gly Thr Leu Phe Ser Arg Ser Val Ile Ala Phe Glu Val Leu Lys Trp
 65 70 75

09847399.05001

gca ggc gcg gct tac ttg att tgg ctg gga atc cag cag tgg cgc gcc 832
Ala Gly Ala Ala Tyr Leu Ile Trp Leu Gly Ile Gln Gln Trp Arg Ala
80 85 90
gct ggt gca att gac ctt aaa tcg ctg gcc tct act caa tcg cgt cga 880
Ala Gly Ala Ile Asp Leu Lys Ser Leu Ala Ser Thr Gln Ser Arg Arg
95 100 105
cat ttg ttc cag cgc gca gtt ttt gtg aat ctc acc aat ccc aaa agt 928
His Leu Phe Gln Arg Ala Val Phe Val Asn Leu Thr Asn Pro Lys Ser
110 115 120
att gtg ttt ctg gcg gcg cta ttt ccg caa ttc atc atg ccg caa cag 976
Ile Val Phe Leu Ala Ala Leu Phe Pro Gln Phe Ile Met Pro Gln Gln
125 130 135 140
ccg caa ctg atg cag tat atc gtg ctc ggc gtc acc act att gtg gtc 1024
Pro Gln Leu Met Gln Tyr Ile Val Leu Gly Val Thr Thr Ile Val Val
145 150 155
gat att att gtg atg atc ggt tac gcc acc ctt gct caa cgg att gct 1072
Asp Ile Ile Val Met Ile Gly Tyr Ala Thr Leu Ala Gln Arg Ile Ala
160 165 170
cta tgg att aaa gga cca aag cag atg aag gcg ctg aat aag att ttc 1120
Leu Trp Ile Lys Gly Pro Lys Gln Met Lys Ala Leu Asn Lys Ile Phe
175 180 185
ggc tcg ttg ttt atg ctg gtg gga gcg ctg tta gca tcg gcg agg cat 1168
Gly Ser Leu Phe Met Leu Val Gly Ala Leu Leu Ala Ser Ala Arg His
190 195 200
gcg tgaaaaataa tgtcggatgc ggcgtaaac 1200
Ala
205

<210> 2

<211> 205

<212> PRT

<213> Escherichia coli

<400> 2

Met Thr Leu Glu Trp Trp Phe Ala Tyr Leu Leu Thr Ser Ile Ile Leu
1 5 10 15
Thr Leu Ser Pro Gly Ser Gly Ala Ile Asn Thr Met Thr Thr Ser Leu
20 25 30
Asn His Gly Tyr Pro Ala Gly Gly Val Tyr Cys Trp Ala Ser Asp Arg
35 40 45
Thr Gly Asp Ser Tyr Cys Ala Gly Trp Arg Gly Val Gly Thr Leu Phe
50 55 60
Ser Arg Ser Val Ile Ala Phe Glu Val Leu Lys Trp Ala Gly Ala Ala
65 70 75 80
Tyr Leu Ile Trp Leu Gly Ile Gln Gln Trp Arg Ala Ala Gly Ala Ile
85 90 95
Asp Leu Lys Ser Leu Ala Ser Thr Gln Ser Arg Arg His Leu Phe Gln
100 105 110
Arg Ala Val Phe Val Asn Leu Thr Asn Pro Lys Ser Ile Val Phe Leu

00047392-050301

115	120	125
Ala Ala Leu Phe Pro Gln Phe Ile Met Pro Gln Gln Pro Gln Leu Met		
130	135	140
Gln Tyr Ile Val Leu Gly Val Thr Thr Ile Val Val Asp Ile Ile Val		
145	150	155
Met Ile Gly Tyr Ala Thr Leu Ala Gln Arg Ile Ala Leu Trp Ile Lys		160
165	170	175
Gly Pro Lys Gln Met Lys Ala Leu Asn Lys Ile Phe Gly Ser Leu Phe		
180	185	190
Met Leu Val Gly Ala Leu Leu Ala Ser Ala Arg His Ala		
195	200	205

0047391 05031